

# Talking with your doctor

Opioid dependence can be treated. Talking with your healthcare team keeps them aware of your situation so they may adjust your treatment plan accordingly.

Speak frankly and honestly. What you tell your doctor is private and should not be shared without your permission.

Add your own questions, and take this guide along on your doctor visit. Use it to help you bring up things you want to discuss. This list is not all inclusive. Your doctor may require additional information from you.

## What does my doctor need to know before I start treatment?

Be specific. Tell your doctor:

- What happened with your previous treatment, including
  - What was working
  - What you were dissatisfied with
- If this is your first time being treated with medication
- What you're looking for from your treatment—your short- and long-term goals
- Anything else that might help your doctor understand your concerns

## Get all of your questions answered

Create a list of questions to ask your doctor, and bring them along so you don't forget anything.

You might be wondering:

- How do my previous treatment and SUBOXONE Film differ?
- What other treatment options are available?
- How can treatment with SUBOXONE Film help me?
- What side effects have people had with SUBOXONE Film?
- How long does treatment typically last?
- Will I experience withdrawal?
- Why is it important to get my dosage right?
- Do I need counseling?
- What factors could put my treatment at risk, and how do I minimize that risk?
- What's my role in treatment?

Write in your own questions here:

1 \_\_\_\_\_


2 \_\_\_\_\_

3 \_\_\_\_\_

4 \_\_\_\_\_

5 \_\_\_\_\_

Please see attached Important Safety Information



# Important Safety Information

## INDICATION

SUBOXONE® (buprenorphine and naloxone) Sublingual Film (CIII) is a prescription medicine indicated for treatment of opioid dependence and should be used as part of a complete treatment plan to include counseling and psychosocial support.

**Treatment should be initiated under the direction of physicians qualified under the Drug Addiction Treatment Act.**

## IMPORTANT SAFETY INFORMATION

Do not take SUBOXONE Film if you are allergic to buprenorphine or naloxone as serious negative effects, including anaphylactic shock, have been reported.

SUBOXONE Film can be abused in a manner similar to other opioids, legal or illicit.

SUBOXONE Film contains buprenorphine, an opioid that can cause physical dependence with chronic use. Physical dependence is not the same as addiction. Your doctor can tell you more about the difference between physical dependence and drug addiction. Do not stop taking SUBOXONE Film suddenly without talking to your doctor. You could become sick with uncomfortable withdrawal symptoms because your body has become used to this medicine.

SUBOXONE Film can cause serious life-threatening breathing problems, overdose and death, particularly when taken by the intravenous (IV) route in combination with benzodiazepines or other medications that act on the nervous system (ie, sedatives, tranquilizers, or alcohol). It is extremely dangerous to take nonprescribed benzodiazepines or other medications that act on the nervous system while taking SUBOXONE Film.

You should not drink alcohol while taking SUBOXONE Film, as this can lead to loss of consciousness or even death.

Death has been reported in those who are not opioid dependent.

Your doctor may monitor liver function before and during treatment.

SUBOXONE Film is not recommended in patients with severe hepatic impairment and may not be appropriate for patients with moderate hepatic impairment. However, SUBOXONE Film may be used with caution for maintenance treatment in patients with moderate hepatic impairment who have initiated treatment on a buprenorphine product without naloxone.

**Please see Full Prescribing Information.**

SUBOXONE® is a registered trademark of Indivior UK Limited

Keep SUBOXONE Film out of the sight and reach of children. Accidental or deliberate ingestion of SUBOXONE Film by a child can cause severe breathing problems and death.

Do not take SUBOXONE Film before the effects of other opioids (eg, heroin, hydrocodone, methadone, morphine, oxycodone) have subsided as you may experience withdrawal symptoms.

Injecting SUBOXONE may cause serious withdrawal symptoms such as pain, cramps, vomiting, diarrhea, anxiety, sleep problems, and cravings.

Before taking SUBOXONE Film, tell your doctor if you are pregnant or plan to become pregnant. If you are pregnant or become pregnant while taking SUBOXONE Film, alert your doctor immediately and you should report it using the contact information provided below.\*

Neonatal withdrawal has been reported following the use of buprenorphine by the mother during pregnancy.

Before taking SUBOXONE Film, talk to your doctor if you are breastfeeding or plan to breastfeed your baby. SUBOXONE can pass into your breast milk. You and your doctor should consider the development and health benefits of breastfeeding along with your clinical need for SUBOXONE Film and should also consider any potential adverse effects on the breastfed child from the drug or from the underlying maternal condition.

Do not drive, operate heavy machinery, or perform any other dangerous activities until you know how SUBOXONE Film affects you. Buprenorphine in SUBOXONE Film can cause drowsiness and slow reaction times during dose-adjustment periods.

Common side effects of SUBOXONE Film include nausea, vomiting, drug withdrawal syndrome, headache, sweating, numb mouth, constipation, painful tongue, redness of the mouth, intoxication (feeling lightheaded or drunk), disturbance in attention, irregular heartbeat, decrease in sleep, blurred vision, back pain, fainting, dizziness, and sleepiness.

This is not a complete list of potential adverse events associated with SUBOXONE Film. Please see [Full Prescribing Information](#) for a complete list.

\*To report negative side effects associated with taking SUBOXONE Film, please call 1-877-782-6966. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

For more information about SUBOXONE Film, SUBOXONE® (buprenorphine and naloxone) Sublingual Tablets (CIII), or SUBUTEX® (buprenorphine) Sublingual Tablets (CIII), please see the respective Full Prescribing Information and Medication Guide at [www.IndiviorREMS.com](http://www.IndiviorREMS.com)



Manufactured for Indivior Inc., Richmond, VA 23235  
Copyright © 2015 Indivior Inc. SBF-UM-P-1162-1409(1) Printed in USA



**Suboxone®** Sublingual  
(buprenorphine and naloxone) **Film**  
2 mg/0.5 mg • 4 mg/1 mg • 8 mg/2 mg • 12 mg/3 mg